

Similac® Special Care®

PREMATURE INFANT
FORMULA
WITH IRON

30 Cal/fl oz



Similac® Special Care® Liqui-Mix™ System*

*For safe** delivery of high-calorie,
nutrient-dense formula!*

- ▶ **SAFE**—Commercially sterile liquid meets ADA and CDC† recommendations to reduce risk of contamination
- ▶ **VERSATILE**—Mixes easily to create a variety of calorically-dense formulas
- ▶ **SIMPLE**—Easy-to-use 1-to-1, 2-to-1 system!

**Upon discharge, recommend
NeoSure® for excellent
catch-up growth during
the first year of life.‡**



* Patent Pending

** Proper hygiene, handling and storage are important when preparing infant formulas. Always follow your hospital's policies and procedures regarding safe handling practices when preparing infant feedings to prevent the possibility of contamination.

† American Dietetic Association, Centers for Disease Control and Prevention

‡ Carver JD, Wu PYK, Hall RT, et al. Growth of preterm infants fed nutrient-enriched or term formula after hospital discharge. Pediatrics. 2001;107:683-689.

© 2008 Abbott Laboratories Inc.
69268.001/NOVEMBER 2008
LITHO IN USA

Abbott Nutrition
Abbott Laboratories
Columbus, OH 43219-3034

 **Abbott**
Nutrition

NUTRIENT COMPOSITION OF SIMILAC® SPECIAL CARE® (SSC) PRODUCTS

PER 100 mL Nutrients	SSC w/Iron 20	SSC w/Iron 24	SSC w/Iron 26 ¹	SSC w/Iron 27 ¹	SSC w/Iron 28 ¹	SSC w/Iron 30
Energy, Cal	68	81	88	91	95	101
Volume, mL	100	100	100	100	100	100
Protein, g	2.03	2.43	2.64	2.74	2.84	3.04
Fat, g	3.67	4.41	5.17	5.56	5.94	6.71
Carbohydrate, g	7.0	8.4	8.2	8.1	8.0	7.8
Calcium, mg	122	146	158	164	170	183
Phosphorus, mg	68	81	88	91	95	101
Magnesium, mg	8.1	9.7	10.6	11.0	11.4	12.2
Iron, mg	1.22	1.46	1.58	1.64	1.70	1.83
Zinc, mg	1.01	1.22	1.32	1.37	1.42	1.52
Manganese, mcg	8.0	10	11	11	11	12
Copper, mcg	169	203	220	228	237	254
Iodine, mcg	4.0	5.0	5.0	5.0	6.0	6.0
Sodium, mg (mEq)	29 (1.3)	35 (1.5)	38 (1.6)	39 (1.7)	41 (1.8)	44 (1.9)
Potassium, mg (mEq)	87 (2.2)	105 (2.7)	113 (2.9)	118 (3.0)	122 (3.1)	131 (3.4)
Chloride, mg (mEq)	55 (1.6)	66 (1.9)	71 (2.0)	74 (2.1)	77 (2.2)	82 (2.3)
Vitamin A, IU	845	1014	1099	1141	1184	1268
Vitamin D, IU	101	122	132	137	142	152
Vitamin E, IU	2.7	3.2	3.5	3.7	3.8	4.1
Vitamin K, mcg	8.1	9.7	10.6	11.0	11.4	12.2
Thiamin (B ₁), mcg	169	203	220	228	237	254
Riboflavin (B ₂), mcg	419	503	545	566	587	629
Vitamin B ₆ , mcg	169	203	220	228	237	254
Vitamin B ₁₂ , mcg	0.37	0.45	0.48	0.50	0.52	0.56
Niacin, mcg	3381	4058	4396	4565	4734	5072
Folic Acid, mcg	25	30	32.5	33.8	35	37.5
Pantothenic Acid, mcg	1285	1542	1670	1735	1799	1927
Biotin, mcg	25	30	32.5	33.8	35	37.5
Vitamin C, mg	25	30	33	34	35	38
Choline, mg	7.0	8.0	9.0	9.0	9.0	10
Inositol, mg	27	32	35	37	38	41
Potential RSL, mOs	18.8	22.6	24.5	25.4	26.3	28.2
Approx Osmolality, mOsm/kg water	235	280	295	305	310	325

¹Can only be made by using the Similac Special Care Liqui-Mix™ System.

Recipes for various formula concentrations using the Similac® Special Care® Liqui-Mix™ System*

To Prepare:

**26 Cal/fl oz
formula**

**27 Cal/fl oz
formula**

**28 Cal/fl oz
formula[†]**

Mix the Following:

*Measure liquid formulas by volume when mixing, as fill weights can vary. Try the Volu-Feed® bottle.

[†]EXAMPLE: To make 1-fl-oz (approx 30 mL) of 28 Cal formula, mix approx. 10mL of SSC w/Iron 24 & 20mL of SSC w/Iron 30.